



Advertisement



AAOS
AMERICAN ACADEMY OF
ORTHOPAEDIC SURGEONS

OrthoInfo

Your connection to expert orthopaedic information

[Find an Orthopaedist](#)

[AAOS.org](#)

- Home
- About Us
- Glossary
- Español
- Videos & Multimedia
- Resources
- For Physicians

Parts of the Body

- Shoulder & Elbow
- Hand & Wrist
- Hip & Thigh
- Knee & Lower Leg
- Foot & Ankle
- Neck & Back

Health Centers

- Broken Bones & Injuries
- Diseases & Conditions
- Arthritis
- Tumors
- Sports Injuries & Prevention
- Children
- Bone Health
- Health & Safety

Treatment

- Treatments & Surgeries
- Joint Replacement
- Rehabilitation Exercise and Conditioning Handouts

Your Healthcare

- Patient Safety
- Patient Stories
- Resources

Dislocated Shoulder

The shoulder joint is the body's most mobile joint. It can turn in many directions. But, this advantage also makes the shoulder an easy joint to dislocate.

A partial dislocation (subluxation) means the head of the upper arm bone (humerus) is partially out of the socket (glenoid). A complete dislocation means it is all the way out of the socket. Both partial and complete dislocation cause pain and unsteadiness in the shoulder.



Shoulder instability

Symptoms

Symptoms to look for include:

- Swelling
- Numbness
- Weakness
- Bruising

Sometimes dislocation may tear ligaments or tendons in the shoulder or damage nerves.

The shoulder joint can dislocate forward, backward, or downward. A common type of shoulder dislocation is when the shoulder slips forward (anterior instability). This means the upper arm bone moved forward and down out of its joint. It may happen when the arm is put in a throwing position.

[Top of page](#)

Diagnosis

The muscles may have spasms from the disruption, and this can make it hurt more. When the shoulder dislocates time and again, there is shoulder instability.

The doctor will examine the shoulder and may order an X-ray. It is important that the doctor know how the dislocation happened and whether the shoulder had ever been dislocated before.

[Top of page](#)

Treatment

The doctor will place the ball of the upper arm bone (humerus) back into the joint socket. This process is called closed reduction. Severe pain stops almost immediately once the shoulder joint is back in place.

[Top of page](#)

Rehabilitation

Your doctor may immobilize the shoulder in a sling or other device for several weeks following treatment. Plenty of rest is needed. The sore area can be iced 3 to 4 times a day.

After the pain and swelling go down, the doctor will prescribe rehabilitation exercises for you. These help restore the shoulder's range of motion and strengthen the muscles. Rehabilitation may also help prevent dislocating the shoulder again in the future. Rehabilitation will begin with gentle muscle toning exercises. Later, weight training can be added.

Like 37

Tweet

Print Article

Related Topics

[Shoulder Trauma \(Fractures and Dislocations\)](#)

If shoulder dislocation becomes a chronic condition, a brace can sometimes help. However, if therapy and bracing fail, surgery may be needed to repair or tighten the torn or stretched ligaments that help hold the joint in place, particularly in young athletes.

[Top of page](#)

Last reviewed: October 2007

AAOS does not endorse any treatments, procedures, products, or physicians referenced herein. This information is provided as an educational service and is not intended to serve as medical advice. Anyone seeking specific orthopaedic advice or assistance should consult his or her orthopaedic surgeon, or locate one in your area through the AAOS ["Find an Orthopaedist"](#) program on this website.

[Editorial Board & Staff](#)

[Contributors](#)

[Online Agreements](#)

[Linking Policy](#)

[Advertising & Sponsorship](#)

[Privacy Policy](#)

[AAOS News Bureau](#)

Copyright ©1995-2015 by the American Academy of Orthopaedic Surgeons. All material on this website is protected by copyright. All rights reserved. This website also contains material copyrighted by third parties.